

Pure artemisinin, or Qinghaosu, is the active constituent of the herb Artemisia annua (sweet wormwood). Good quality Artemisia annua contains 0.3-0.5% artemisinin, so pure artemisinin provides hundreds of times more of the active constituent artemisinin than the whole herb itself. Research has shown artemisinin to be particularly beneficial in balancing the microbiology of the GI tract.<sup>\*</sup>

Our **Artemisinin** has had independent cell tests verifying its effectiveness, and we do independent potency assays on every batch, using HPLC. Numerous practitioners have chosen NutriCology<sup>®</sup> artemisinin as part of the nutritional supplement program they utilize in their practice.<sup>\*</sup>

PhytoArtemisinin provides artemisinin with gleditsia extract, a phytosaponin.



along with herbal extracts, such as berberine, grapefruit seed extract and oregano oil." - Leo Galland, MD

"Artemisinin is a powerful oxidant. I have used it orally

"I have been very impressed by the efforts of NutriCology<sup>®</sup> to perform both purity analysis and activity analysis on its artemisinin. For me, it was not even necessary, since I had already observed the awesome clinical efficacy of the product in patient after patient. I am grateful that this product is available and at a most reasonable price." - Robert Jay Rowen, MD

#52160 • Artemisinin 90 vegetarian capsules

#55680 • Artemisinin 300 vegetarian capsules



800.545.9960 info@nutricology.com www.nutricology.com



## Artemisinin

90 vegetarian capsules • 52160 300 vegetarian capsules • 55680

Supplement Fac	ts
Serving Size Servings Per Contair	45 or 150 ner 45
Amount Per Serving % Daily Valu	
Amount Per Serving	% Daily Value
Amount Per Serving Artemisinin	% Daily Value           200 mg         †



Other ingredients: Hydroxypropyl methylcellulose, microcrystalline cellulose, rice hull concentrate, gum arabic, sunflower oil.

**Suggested Use:** As a dietary supplement, 1 or 2 capsules one or two times daily before meals, or as directed by a healthcare practitioner. Artemisinin is best taken as part of a program that includes appropriate liver support nutrients. Sensitive individuals may want to take with food.

## PhytoArtemisinin

90 vegetarian capsules • 54990

Supplement Facts		
Serving Size Servings Per Container	2 Capsu	les 45
Amount Per Serving	% Daily Val	ue
Gleditsia (Spine) Extract (30%	Saponins)	
Gleditsia (Spine) Extract (30%	Saponins) 700 mg	†
Gleditsia (Spine) Extract (30%	' '	† †



Other ingredients: Hydroxypropyl methylcellulose, L-leucine.

**Suggested Use:** As a dietary supplement, 1 or 2 capsules one or two times daily before meals, or as directed by a healthcare practitioner. Artemisinin is best taken as part of a program that includes appropriate liver support nutrients. Sensitive individuals may want to take with food.

**Caution:** Not indicated for pregnant or nursing women. These products should only be used under the guidance of a qualified healthcare practitioner with ongoing monitoring of liver enzymes and hemoglobin during their use. In rare cases may cause idiosyncratic liver dysfunction. Combining with antioxidants or iron may theoretically decrease effectiveness.

References:
Ames.JR, Ryan MD, et al. J.: Free Rad Biol. Med 1985, 1:353-61.
Beekman AC, Wierenga PK, Woerdenbag HJ, Van Uden W, Pras N, Konings AW, Wikstrom HV. J Pharm Pharmacol. 1997 Dec;49(12):1254-8.
Bharel S, Claiki M, Abdim F, Shrivastava S, Ficteripai Vol LSWI MD, 52, 2013.
China Cooperative Research Group. J. Trad. Chin. Med 2, 17, 1982.
Efferth T, Dunnar H, Sauedrory A, Miyachi H, Chinambar CR. Oncol. 2001. Apr:18(4):757-73.
Fiefrich T, Dunnar H, Sauedrory A, Miyachi H, Chinambar CR. Oncol. 2001. Apr:18(4):767-73.
Fiefrich T, Dunnar H, Sauedrory A, Molachi H, Chinambar CR. Oncol. 2001. Apr:18(4):767-73.
Fiefrich T, Dunnar S, Srivastava M, Abdim Z, Tichreanja Vol LSWI 11No 5, 1979.
Keith Amold, Tran Tinh Hien, et al. Transactions of the Royal Society of Tropical Medicine and Hygiene (1990) 84:499-502.
Kayaman D, Science, 1985, Vol. 238. May 31, p. 1049.
Krughard SK, Yuthavong, Trans. R Soc. Trop. Med Hyg 1987, 81:710-4.
Lee CH, Hong H, Shin J, Yoan J, Lee W. Biochem Biophys Res Commun. 2000 Aug 2;274(2):359-69.
Levander OA, Ager AJ, Morris VC. Am. J. Clin. Nutr. 1989; 5:346-32.
Lu Zhonggu Y, Davad Z, May 213, 1055-60.
McHanna MG, Ward SA, Med Trop (Mars). 1998; 5:846-32.
McLean WG, Ward SA, Med Trop (Mars). 1998; 5:846-32.
Muchangamang S, Widerston M, Naik YS.
Mannenk B, Haaler JA, Int J Cancer. 2002 Leck 21(3):1655-61.
McHanna MG, Ward SA, Med Trop (Mars). 1998; 5:845-82.
McLean WG, Ward SA, Med Trop (Mars). 1998; 5:845-82.
McLean WG, Ward SA, Med Trop (Mars). 1998; 5:89(3):2013.
Mohangamag S, Widerston M, Naik YS.
Mannenet B, Haaler JA, Int J Cancer. 2002 Peb 1097(5):700-5.
Phan GT, de Vreis PJ, Tran BQ, Le HQ, Nguyen TV, Heiterkamp SH, Kager PA. Trop Med Int Health. 2002 Oct;7(10):858-64.
Posner GH, Paik H, Burston K, Cholan P, Keneler TW, Xie S, Shapiro TA. Bind Orde. C

NutriCology® | 2300 South Main Street, South Salt Lake, UT 84115 | 800.545.9960 | info@nutricology.com | www.nutricology.com